

BMJ Open Which cuff should I use? Indirect blood pressure measurement for the diagnosis of hypertension in patients with obesity: a diagnostic accuracy review

Greg Irving,¹ John Holden,² Richard Stevens,³ Richard J McManus³

To cite: Irving G, Holden J, Stevens R, *et al*. Which cuff should I use? Indirect blood pressure measurement for the diagnosis of hypertension in patients with obesity: a diagnostic accuracy review. *BMJ Open* 2016;**6**:e012429. doi:10.1136/bmjopen-2016-012429

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2016-012429>).

Received 25 April 2016
Accepted 10 June 2016



CrossMark

For numbered affiliations see end of article.

Correspondence to

Dr Greg Irving; gi226@cam.ac.uk

ABSTRACT

Objective: To determine the diagnostic accuracy of different methods of blood pressure (BP) measurement compared with reference standards for the diagnosis of hypertension in patients with obesity with a large arm circumference.

Design: Systematic review with meta-analysis with hierarchical summary receiver operating characteristic models. Bland-Altman analyses where individual patient data were available. Methodological quality appraised using Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS2) criteria.

Data sources: MEDLINE, EMBASE, Cochrane, DARE, Medion and Trip databases were searched.

Eligibility criteria: Cross-sectional, randomised and cohort studies of diagnostic test accuracy that compared any non-invasive BP tests (upper arm, forearm, wrist, finger) with an appropriate reference standard (invasive BP, correctly fitting upper arm cuff, ambulatory BP monitoring) in primary care were included.

Results: 4037 potentially relevant papers were identified. 20 studies involving 26 different comparisons met the inclusion criteria. Individual patient data were available from 4 studies. No studies satisfied all QUADAS2 criteria. Compared with the reference test of invasive BP, a correctly fitting upper arm BP cuff had a sensitivity of 0.87 (0.79 to 0.93) and a specificity of 0.85 (0.64 to 0.95); insufficient evidence was available for other comparisons to invasive BP. Compared with the reference test of a correctly fitting upper arm cuff, BP measurement at the wrist had a sensitivity of 0.92 (0.64 to 0.99) and a specificity of 0.92 (0.85 to 0.87). Measurement with an incorrectly fitting standard cuff had a sensitivity of 0.73 (0.67 to 0.78) and a specificity of 0.76 (0.69 to 0.82). Measurement at the forearm had a sensitivity of 0.84 (0.71 to 0.92) and a specificity 0.75 of (0.66 to 0.83). Bland-Altman analysis of individual patient data from 3 studies comparing wrist and upper arm BP showed a mean difference of 0.46 mm Hg for systolic BP measurement and 2.2 mm Hg for diastolic BP measurement.

Conclusions: BP measurement with a correctly fitting upper arm cuff is sufficiently sensitive and specific to diagnose hypertension in patients with obesity with a large upper arm circumference. If a correctly fitting upper arm cuff cannot be applied, an incorrectly fitting standard

Strengths and limitations of this study

- Study quality was assessed using Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS2).
- Forest plots of sensitivity and specificity were created for each study and estimates of sensitivity and specificity in receiver operating characteristic space for each comparison were plotted.
- Individual patient data were obtained and the Bland-Altman method used to plot differences in measurement against reference standard.
- Insufficient evidence was available to compare forearm, wrist, finger blood pressure measurement with invasive blood pressure.
- All included studies used body mass index (BMI) >30 as an indicator of obesity. Although BMI generally increases as adiposity increases, due to variation in body composition, it is not the most accurate measure of body adiposity.

size cuff should not be used and BP measurement at the wrist should be considered.

INTRODUCTION

Approximately 671 million individuals worldwide are obese and subsequently mean arm circumference has increased.^{1 2} As a result, healthcare professionals are increasingly faced with situations where blood pressure (BP) measurement with a standard (or even large sized upper arm cuff) is not possible.² This is important because errors in BP measurement are greater when the cuff used is too small relative to the patient's arm circumference.³ In the USA, over 30% of the population have a large arm circumference as a consequence of obesity, a figure which can rise to over 60% in some clinics.^{4 5}

Standard cuffs are ~22–32 cm with large cuffs typically ranging from 32 to 42 cm.⁶

Where available cuffs are not large enough for an individual's upper arm, clinicians may use a variety of different methods for BP measurement including a correctly fitting extra large cuff, forearm BP, wrist BP, finger BP or ambulatory BP.⁷ However, there is considerable uncertainty as to the diagnostic accuracy of these different approaches for obese people and hence the optimum alternative test.² Currently, all of the major hypertension guidelines recommend different approaches (table 1).

Understanding which alternative method to use is important because hypertension is common in patients with obesity and the consequences of missing the diagnosis is potentially severe at the individual and population level.¹¹ This review aimed to determine the diagnostic accuracy of upper arm, forearm, wrist and finger BP measurement compared with the reference standards of invasive arterial, upper arm (correctly fitting cuff) and/or ambulatory BP measurement (ABPM) for the diagnosis of hypertension in patients with obesity with a large arm circumference.

METHODS

Search methods for identification of studies

MEDLINE, EMBASE, Cochrane database, DARE, Medion and the TRIP database were searched. No language or publication status restrictions were applied. To increase the sensitivity of the search, methodology filters were not used as these have been found to miss relevant studies when searching for diagnostic accuracy studies.¹² Reference lists from all primary studies and reviews identified were hand searched.

Inclusion criteria

Studies comparing upper arm, forearm, wrist or finger BP with reference standards of intra-arterial, upper arm (correctly fitting cuff) or ABPM for the diagnosis of hypertension in adult (over 18 years) patients with obesity were included. For each index test, all cuff shapes and sizes were considered. Thresholds for hypertension of 140/90 mm Hg (clinic and intra-arterial measurement) and 135/85 mm Hg (ABPM) were used

Table 1 Implications for policy

Guideline	Guidance	Comment
American Heart Association ⁸	"In patients with morbid obesity, one will encounter very large arm circumferences...In this circumstance, the clinician may measure blood pressure from a cuff placed on the forearm and listening for sounds over the radial artery or use a validated wrist blood pressure monitor held at the level of the heart."	There was insufficient evidence to undertake subgroup analyses for patients with an arm circumference >50 cm. However, on the basis of the available evidence in this review, if a correctly fitting upper arm cuff cannot be placed then measurement at the wrist is preferable to measurement at the forearm.
European Society of Cardiology ⁹	"Devices worn on the wrist are currently not recommended but their use might be justified in obese subjects with extremely large arm circumference."	This review would support the use of wrist BP devices in patients with obesity if a correctly fitting upper arm cuff cannot be applied and the device was used at the level of the heart.
British Hypertension Society/ National Institute for Clinical Excellence ¹	"A large cuff is still too small. What should I do? Contact the manufacturer of the blood pressure monitor. They may be able to supply an extra large cuff." No specific guidance is offered for the use of wrist BP measurement in patients with obesity with a large arm circumference. "The bladder of the cuff should fit around at least 80% of the arm but not more than 100%. A cuff that does not fit properly will not give an accurate reading so it is important to use the right size."	Compared with the reference test of invasive BP, a correctly fitting upper arm BP cuff is sufficiently sensitive and specific for the diagnosis of hypertension in patients with obesity. This holds true for patients with BMI>35. It was not possible to undertake a subgroup analysis of those with arm circumference >40 or 50 cm. There was good evidence that an incorrectly fitting standard cuff was not as accurate as a correctly fitting cuff. Both wrist and forearm BP measurements were more accurate than an incorrectly fitting upper arm cuff.
WHO/International Society for Hypertension ¹⁰	No specific guidance is offered for the use of wrist BP measurement in patients with obesity with a large arm circumference.	Compared with the reference test of invasive BP, a correctly fitting upper arm BP cuff is sufficiently sensitive and specific for the diagnosis of hypertension in patients with obesity. This holds true for patients with BMI>35. It was not possible to undertake a subgroup analysis of those with arm circumference >40 or 50 cm.

BMI, body mass index; BP, blood pressure.

and both manual and automated BP measurements were considered. Obese adults were defined by either an upper arm circumference ≥ 35 cm, a body mass index (BMI) ≥ 30 or by direct measurement of percentage body fat ($\geq 25\%$ for men and $\geq 30\%$ for women). Prespecified subgroups included those with arm circumference ≥ 40 cm or ≥ 50 cm or BMI ≥ 35 .

Included study designs were diagnostic cross-sectional, randomised and cohort studies. Studies were excluded if participants were: receiving antihypertensive treatment at the time of comparison or pregnant or a hospital inpatient. Studies from which data could not be extracted were included in the descriptive part of the review but excluded from subsequent analyses.

Selection and data extraction

To determine inclusion or exclusion of each potential study, GI and JH independently reviewed the results of the search by title/abstract and when required by the full text of the study. The resulting list of citations was then reviewed for inclusion and a third author (RJM) arbitrated the final selection decisions.

A standardised data extraction form was used to identify study characteristics and results for each included publication which were independently extracted by GI and JH. Study quality was assessed using Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS2).¹³ Where possible, data were extracted into 2x2 contingency tables. All authors were contacted directly if insufficient data were available in the published report of a given study and to request individual patient data.

Statistical analysis and data synthesis

Forest plots of sensitivity and specificity were created for each study and estimates of sensitivity and specificity in receiver operating characteristic (ROC) space for each comparison were plotted. The diagnostic performance of each index test was ascertained and heterogeneity in such performance estimated (see below). The Metandi and Midas procedures in STATA V.13.0 were used to fit the hierarchical summary receiver operating curve (HSROC) models (R Harbord. METANDI: Stata module to perform meta-analysis of diagnostic accuracy. Statistical Software Components 2008; Stata Corporation. Stata Statistical Software Release V.13.0: Programming; Stata Corporation, 2001). Where there was a sufficient number of studies ($n \geq 5$) the HSROC model was used to derive inferences about diagnostic test accuracy and heterogeneity in test performance including the summary curve.¹⁴ Where individual patient data were available, the Bland-Altman method was used to plot differences in measurement against reference standard.¹⁵

Investigations of heterogeneity and sensitivity analyses

Heterogeneity was evaluated by visual inspection of Forest plots of sensitivity and specificity, ROC plots of the data and using the χ^2 test. The influence of

differences in the test characteristics (manual vs automated BP device), study population (primary care vs hospital outpatient) and methodological quality (scores on items of the QUADAS2 checklist) was investigated where it was appropriate to do so.

Where there was evidence of differences between studies sensitivity analyses were undertaken based on:

1. Measurement of obesity: indirect/direct methods, for example, bioelectrical impedance analysis (BIA) or skin fold thickness.
2. Risk of bias according to QUADAS2 criteria: low/high risk of bias.
3. Year of study: $<25/\geq 25$ years old.

Reporting bias was assessed using the test for funnel plot asymmetry.¹⁶

Patient involvement

The design of the review was informed by a search of patient uncertainties in the UK Database of Uncertainties of the Effects of Treatment (DUETS) and patient research priorities identified by the James Lind Alliance.

RESULTS

Results of the search

A total of 4037 studies were identified (excluding duplicates). The full text of 164 papers was reviewed for eligibility (figure 1) and 37 studies were included (32 published in English). One was a randomised trial and one a case-control study;^{17 18} the remainder were cross-sectional studies. Twenty studies (26 comparisons) had extractable data, all of which were of cross-sectional design. There were no disagreements between authors in relation to the number of studies eligible for inclusion ($\kappa=1.0$). Individual patient data were available from four authors.

Methodological quality of included studies

Quality assessment of the 20 included studies and the 16 studies that could not contribute data, found no studies that satisfied all the QUADAS2 criteria as all had some degree of methodological weakness and/or lacked reporting clarity (figure 2). Blinding of reference tests, blinding of index tests and acceptable delay between tests were particularly poorly reported.

Findings

Six studies were found which used invasively measured BP as the reference standard and a further 17 using a properly fitting upper arm cuff. One study used ABPM as a reference standard. Details of included studies are shown in table 2 and results are presented by reference test below.

Reference test: invasively measured BP

Six studies were found with extractable data (eight data sets) comparing upper arm BP with invasive




Figure 1 *Date of search 1/6/2015. PRISMA flow diagram. DTA, diagnostic test accuracy.

measurement in obesity yielding a pooled sensitivity of 0.87 (0.79 to 0.93) and a specificity of 0.85 (0.64 to 0.95).^{18 27 29 32–34} Only three studies were inside the 95% CIs for the summary receiver operating curve (SROC) summary point. The five outlying studies had a relatively small sample size (5–20 participants). Deeks funnel plot asymmetry test had a non-significant p value ($p=0.53$) for the slope coefficient suggesting no evidence of publication bias. Only one study was included for the subgroup analysis of those patients with BMI>35.²⁷ This resulted in a sensitivity of 0.88 (0.47 to 1.00) and a specificity of 0.71 (0.40 to 0.90). There were insufficient data to perform any other subgroup analyses. Across the six studies there was no difference in reference threshold and all avoided verification bias.

	Risk of Bias				Applicability Concerns		
	Patient Selection	Index Test	Reference Standard	Flow and Timing	Patient Selection	Index Test	Reference Standard
Bennett 1987	?	?	?	?	?	+	+
Berntsen 1998	?	?	?	?	+	+	?
Bertrand 1956	+	+	+	+	?	+	+
Blackburn 1965	?	+	?	?	?	?	?
Bovet 1994	+	?	?	?	+	?	+
Chiolero 2006	?	?	?	?	+	?	?
Cuckson 2004	?	?	?	+	+	?	+
de Senarclens 2008	?	?	?	+	+	+	?
Domiano 2008	+	?	?	+	+	+	+
Doshi 2010	+	?	?	+	+	?	+
Genc 2008	?	?	?	?	?	+	+
Guagnano 1994	?	?	?	+	+	+	+
Guagnano 1996	+	+	+	+	+	+	+
Guagnano 1998	+	?	?	+	+	+	+
Julien 1988	?	?	?	+	+	+	+
King 1967	?	?	?	?	?	?	+
Kotsis 2005	?	?	?	?	+	+	+
Kvols 1969	?	?	?	?	+	?	+
Leblanc 2013	?	?	+	+	+	+	+
Linforts 1984	?	+	+	?	+	+	?
Maxwell 1982	?	?	?	?	+	?	+
Maxwell 1985	?	?	+	+	+	?	+
Nielsen 1983	+	?	?	+	+	+	+
Palatini 2012	?	+	+	+	+	+	+
Pang 2006	+	+	+	+	?	+	+
Pierin 2004	?	?	?	+	+	+	+
Poncelet 1992	+	?	?	+	?	?	+
Schell 2005	?	?	?	+	+	?	+
Schell 2006	?	?	?	?	+	?	+
Simpson 1965a	+	?	+	+	+	?	+
Stergiou, 2008	?	?	?	+	+	+	+
Stolt 1990a	+	?	?	+	+	+	+
Stolt 1993	?	?	?	+	+	+	+
Umana 2006	+	?	?	+	+	?	+
van Montfrans 1987	+	?	?	?	+	+	+
Vinyoles 2005	+	?	?	+	+	+	+
Warembourg 1987	+	?	?	+	+	+	+

High ? Unclear Low

Figure 2 Risk of bias and applicability concerns summary with review authors' judgements for each included study.

Table 2 Characteristics of included studies

Study	Sample size (patients with obesity)	Mean age	Male (%)	Study population	Study design
Bennett <i>et al</i> ^{19*}	26	–	–	Community	Cross-sectional
Berntsen <i>et al</i> ^{20*}	4	65	–	Community	Cross-sectional
Bertrand <i>et al</i> ^{18*}	16	–	–	Community	Case-control
Blackburn 1965 ³⁷	76	–	100	Hospital outpatient	Cross-sectional
Bovet 1994 ³⁸	103	–	43.7	Hospital outpatient	Cross-sectional
Chiolero <i>et al</i> ^{21*}	62	–	49.7	Community	Cross-sectional
Cuckson <i>et al</i> ^{22*}	20	–	–	Hospital outpatients	Cross-sectional
de Senarclens <i>et al</i> ^{23*}	12	39.9	27	Hospital outpatient	Cross-sectional
Domiano <i>et al</i> ^{24*}	42	50.7	56	Community	Cross-sectional
Doshi <i>et al</i> ^{3*}	103	39.9	27	Hospital outpatient	Cross-sectional
Genc <i>et al</i> ³⁹	32	51.4	44.8	Hospital outpatient	Cross-sectional
Guagnano <i>et al</i> ⁴⁰	97	48.7	35	Hospital outpatient	Cross-sectional
Guagnano <i>et al</i> ²⁵	94	42.9	0	Hospital outpatient	Cross-sectional
Guagnano <i>et al</i> ^{26*}	339	41	0	Hospital outpatient	Cross-sectional
Julien 1988 ⁴¹	19	51	5	Hospital outpatient	Cross-sectional
King 1967 ⁴²	–	–	–	Hospital outpatient	Cross-sectional
Kotsis 2005 ⁴³	825	53.5	53	Hospital outpatient	Cross-sectional
Kvols 1969 ⁴⁴	–	–	–	Hospital outpatient	Cross-sectional
Leblanc <i>et al</i> ^{27*}	25	–	–	Hospital outpatient	Cross-sectional
Linfors <i>et al</i> ^{28*}	116	51	29	Hospital outpatient	Cross-sectional
Maxwell 1982 ⁴⁵	–	–	–	Hospital outpatient	Cross-sectional
Maxwell 1985 ⁴⁶	–	–	51	Hospital outpatient	Cross-sectional
Nielsen <i>et al</i> ^{29*}	57	41	53	Hospital outpatient	Cross-sectional
Palatini <i>et al</i> ²	–	54	50	Hospital outpatient	Cross-sectional
Pang 2006 ⁴⁷	–	–	–	Hospital outpatient	Cross-sectional
Pierin <i>et al</i> ^{30*}	155	45	10	Hospital outpatient	Cross-sectional
Poncelet <i>et al</i> ^{31*}	6	52	33	Hospital outpatient	Cross-sectional
Schell 2005 ⁴⁸	204	36.5	51.5	Hospital outpatient	Cross-sectional
Schell 2006 ^{49*}	79	–	46.6	Hospital outpatient	Cross-sectional
Simpson <i>et al</i> ³²	5	–	–	Hospital outpatient	Cross-sectional
Stergiou <i>et al</i> ^{17*}	20	–	–	Community	Randomised trial
Stolt <i>et al</i> ^{33*}	10	–	–	Community	Cross-sectional
Stolt <i>et al</i> ^{34*}	20	48	35	Community	Cross-sectional
Umana <i>et al</i> ⁵⁰	61	62–	62	Hospital outpatient	Cross-sectional
van Montfrans 1987 ⁵¹	19	–	–	Hospital outpatient	Cross-sectional
Vinyoles <i>et al</i> ^{35*}	108	60.2	22.2	Community	Cross-sectional
Warembourg <i>et al</i> ⁵²	10	54	0	Hospital outpatient	Cross-sectional

*Studies with extractable data.

All studies used body mass index ≥ 30 as definition of obesity.

There was no significant evidence of heterogeneity across the six studies (χ^2 : $Q=1.51$, $df=2$, $p=0.24$). Exclusion of the two studies that were >25 years old (1953 and 1965) and used older equipment resulted in a pooled sensitivity of 0.86 (0.77 to 0.92) and a specificity of 0.90 (0.70 to 0.92) and did not influence heterogeneity.^{22 30} The number of studies was too small to calculate a pooled analysis for other a priori stated potential sources of heterogeneity.

No studies with extractable data regarding forearm, wrist or finger BP in obese individuals compared with invasively measured BP were found.

Reference test: correctly fitting upper arm cuff


Sixteen studies (18 comparisons) were found with a reference standard of correctly fitting upper arm cuff

compared with a standard cuff, forearm, wrist or finger BP measurement.

Upper arm BP

Five studies (six data sets) compared a standard sized (ie, too small) upper arm cuff with a correctly fitting upper arm cuff and found a pooled sensitivity of 0.73 (0.67 to 0.78) and a specificity of 0.76 (0.69 to 0.82).^{19–21 26 28} The majority of the studies were within the 95% CI of the SROC summary point. The two outliers had a relatively small sample size in comparison to the other studies ($n=4$ and $n=31$) and the Deeks funnel plot suggested no evidence of publication bias ($p=0.34$). Only one study was included in the subgroup analysis of those patients with $BMI>35$ with resultant wide CIs: sensitivity 1.00 (0.59 to 1.00) and

Figure 3 Summary ROC points with 95% CIs for: upper arm versus correct cuff; forearm BP versus correct cuff; four wrist BP versus correct cuff; five finger versus correct cuff. BP, blood pressure; ROC, receiver operating characteristic.



specificity 0.67 (0.22 to 0.96).²¹ There were insufficient data to perform any other subgroup analyses.

Across the five studies, there was no difference in reference threshold and all studies avoided verification bias. There was no significant statistical evidence of heterogeneity across the six comparisons (χ^2 : $Q=1.62$, $df=2$, $p=0.22$) but there were insufficient data to carry out a pooled analysis for other potential sources of heterogeneity.

Forearm BP

Six studies compared BP measurement at the forearm at the level of the heart with a correctly fitting upper arm cuff and found a sensitivity of 0.84 (0.71 to 0.92) and a specificity of 0.75 (0.66 to 0.83).^{18 24 27 30 35} Four of the studies were within the 95% CI of the SROC summary point. Deeks funnel plot asymmetry test had a non-significant p value ($p=0.50$) for the slope coefficient suggesting no evidence of publication bias. There was statistical evidence of heterogeneity across the five studies (χ^2 : $Q=8.382$, $df=2.00$, $p=0.008$) which appeared largely due to two studies with much earlier publication dates.^{18 32} Excluding these studies resulted in a pooled sensitivity of 0.86 (0.77 to 0.92) and a specificity of 0.90 (0.70 to 0.92) and eliminated heterogeneity (χ^2 : $Q=0.68$, $df=2.00$, $p=0.36$). Further analyses for heterogeneity or subgroups were not possible.

Wrist BP

Five studies considered BP measured at the wrist held at the level of the heart compared with a properly fitting upper arm cuff yielding a pooled sensitivity of 0.92 (0.64 to 0.99) and a specificity of 0.92 (0.85 to 0.97).^{3 17 21 23 31} Funnel plot and χ^2 tests suggested a low probability of publication bias ($p=0.89$) or heterogeneity (χ^2 : $Q=4.26$, $df=2.00$, $p=0.06$), and subgroup analyses were not possible.

Finger BP

One study with two extractable data sets compared finger BP measurement with that from a correctly fitted upper arm cuff and reported a similar specificity (0.57 and 0.61) but a markedly different sensitivity (0.74 and 0.91) from the two comparisons.³¹ With so few data, it was not possible to calculate a pooled summary estimate, assess for heterogeneity or consider subgroups or publication bias.

Comparing methods

A summary ROC plot of all index tests compared with the reference test of a correctly fitting upper arm cuff is provided in figure 3. Visual inspection suggests that measurement at the wrist held at the level of the heart performs best in comparison to the reference.

Table 3 Bland-Altman mean difference values for individual studies of wrist versus correct cuff⁵³

Study	Specificity	Systolic BP Bland-Altman mean difference (limits of agreement \pm 2 SD)	Diastolic BP Bland-Altman mean difference (limits of agreement \pm 2 SD)
Chiolero <i>et al</i> ²¹	0.93	0.51 (−22.7 to 23.7)	1.96 (−16.3 to 20.2)
Doshi <i>et al</i> ⁸	0.91	2.61 (−20.9 to 15.7)	4.98 (−18.1 to 8.2)
de Senarclens <i>et al</i> ²³	0.91	−6.09 (−22.3 to 10.2)	1.36 (−12.6 to 15.3)
Stergiou <i>et al</i> ¹⁷	0.88	4.05 (−14.8 to 22.9)	3.61 (−6.0 to 13.2)
All*		0.46 (−22.7 to 22.6)	2.2 (−14.1 to 18.6)

*Doshi *et al*⁸ not included in pooled result as no individual patient data available. The figure presented is taken from the Bland-Altman analysis provided in the paper.
BP, blood pressure.

Individual patient data

Bland-Altman analysis of the available individual patient data for three studies comparing wrist and upper arm BP measurement with a correctly fitting cuff showed a mean difference of 0.51 mm Hg (limits of agreement −21.7 to 22.6 mm Hg; systolic BP) and 1.96 mm Hg (limits of agreement −14.1 to 18.6 mm Hg) (diastolic BP; table 3).^{17 21 23}

Reference test: ambulatory BP

One study with extractable data compared upper arm BP measurement with that of ambulatory BP in obesity.²⁵ This showed a sensitivity of 0.45 (0.29 to 0.62) and a specificity of 0.71 (0.60 to 0.81). Given that only a single study was identified, we were not able to assess for statistical heterogeneity or undertake subgroup analyses. No studies with extractable data regarding forearm, wrist or finger BP in obese individuals compared with ambulatory monitoring were found.

DISCUSSION

Summary of main results

This review has shown that measurement of BP in an obese individual using a correctly fitting upper arm BP cuff compared with the reference standard of direct arterial measurement provides similar results to those obtained in non-obese adults: sensitivity (0.87 (0.79 to 0.93; obese))

versus (0.67 (0.30 to 0.93; non-obese)) and specificity (0.85 (0.64 to 0.95; obese) versus (0.95 (0.83 to 0.99; non-obese); table 4).³⁴ There is insufficient evidence to comment on how forearm, wrist and finger BP testing perform in relation to invasively measured BP in obesity.

Although CIs around the point estimates overlapped, inspection of SROC curves suggested BP measurement at the wrist is the alternative of choice should a correctly fitting upper arm cuff not be available. Bland-Altman analysis of available individual patient data demonstrated good agreement between wrist and upper BP measurement with a correctly fitting cuff for systolic (mean difference 0.51 mm Hg) and diastolic BP (mean difference 1.96 mm Hg). This falls within the ± 3 mm Hg (British Hypertension Society (BHS) standard) and ± 5 mm Hg (clinically relevant difference) margin of error. However, it is important to note that this analysis was based on a relative small amount of data with wide limits of agreement and that care was taken to minimise the influence of arm–heart hydrostatic pressures by asking the patient to hold the wrist devices at the level of the heart. If this is not performed, then systematic errors in measurement are likely to occur.⁶

Limitations of the review

For a number of the studies included in this review, data were either missing or could not be extracted directly from the published paper in order to construct the

Table 4 Summary of results

Reference	Index	Number of Studies (number of participants)	Sensitivity	Specificity	Positive likelihood ratio	Negative likelihood ratio
Invasive	Correctly fitting cuff	6 (163)	0.87 (0.79 to 0.93)	0.85 (0.64 to 0.95)	3.33 (1.89 to 5.88)	0.18 (0.09 to 0.35)
Correctly fitting cuff	Standard cuff	6 (621)	0.73 (0.67 to 0.78)	0.76 (0.69 to 0.82)	2.74 (2.03 to 3.70)	0.38 (0.30 to 0.47)
Correctly fitting cuff	Forearm cuff	6 (425)	0.84 (0.71 to 0.92)	0.75 (0.66 to 0.83)	3.64 (2.47 to 5.37)	0.17 (0.08 to 0.34)
Correctly fitting cuff	Wrist cuff	5 (217)	0.92 (0.64 to 0.99)	0.92 (0.85 to 0.87)	12.62 (5.90 to 27.0)	0.82 (0.01 to 0.49)

required contingency tables. Authors were contacted directly to obtain these data, and individual patient data made available by three researchers was used to calculate Bland-Altman analyses.

All included studies used BMI >30 as an indicator of obesity which has limitations. Although BMI generally increases as adiposity increases, due to variation in body composition, it is not the most accurate measure of body adiposity.¹ Those with a high proportion of muscle mass could have a high BMI. In a similar way arm circumference as a measure of arm obesity may be confounded in the case of an athlete with a low percentage body fat and large, muscular arms. No studies used direct methods such as BIA, skin fold thickness, underwater weighing. In some populations, the percentage of patients without obesity with a large arm circumference arms can be ~10% (Latman 2013, personal communication).

Some statistical heterogeneity was apparent (forearm vs correctly fitting upper arm cuff) and this could not be explained by reference threshold differences or partial verification. Instead, other factors are likely to have contributed, in particular publication date, suggesting the quality of equipment used may be important. However, it is possible other sources may have contributed including population characteristics (prevalence of hypertension) or test application (operator variability). Where heterogeneity was not detected, it is important to be aware of the limitations of tests for heterogeneity in diagnostic accuracy studies particularly when the number of included studies is small.³⁶

Comparison with the literature

This review is the first, to the best of our knowledge, to summarise the international literature on indirect BP measurement in obesity. This is perhaps reflected in the variation between international guidelines in their recommendations for BP measurement in obesity.^{1 8–10} The American Heart Association currently advocate the use of forearm BP, the European Society of Cardiology advocate wrist BP measurement and the British Hypertension Society suggest contacting manufacturers in order to obtain an extra large upper arm cuff.^{1 8 9} The International Society of Hypertension offers no specific advice.¹⁰

Clinical and policy implications

On the basis of the review, compared with the reference test of invasive BP, a correctly fitting upper arm BP cuff is sufficiently sensitive and specific for the diagnosis of hypertension in patients with obesity. This also hold true for patients with a BMI >35. If a correctly fitting cuff cannot be fitted or is unavailable, then wrist BP measurement appears to be the next best alternative. BP measurement with an incorrectly fitting standard size cuff on the upper arm should be avoided as it is an insufficiently sensitive or specific test for a diagnosis of hypertension.

All included studies had some degree of methodological weakness and/or lacked of clarity in their

reporting. Blinding of reference tests, blinding of index tests and acceptable delay between tests were particularly poorly reported. For this reasons, it is important for any future studies to address these important methodological considerations and follow the QUADAS2 guidance.¹³ On the basis of the paucity of individual patient data, there is now a need for a large diagnostic accuracy study of high methodological quality comparing wrist/forearm BP measurement with upper arm BP. This should aim to include patients with an arm circumference at least 40 cm. To determine percentage arm adiposity more accurately direct methods such as BIA and skin fold thickness could be employed.

CONCLUSIONS

This review set out to identify the diagnostic accuracy of non-invasive BP measurement techniques (upper arm, forearm wrist, finger) for the diagnosis of hypertension in patients with obesity with a large arm circumference compared with three different reference standards: invasive BP, correctly fitting upper arm cuff and ABPM. In conclusion, in the absence of a correctly fitted upper arm cuff, measurement at the wrist appears appropriate, provided that the cuff is held at the level of the heart.

Author affiliations

¹Department of Primary Care, Institute of Public Health and Primary Care, University of Cambridge, Cambridge, UK

²Department of Garswood Surgery, Garswood, UK

³Nuffield Department of Primary Care Health Sciences, NIHR School for Primary Care Research, University of Oxford, Oxford, UK

Acknowledgements The authors would like to thank those authors who shared study data (Bennett, Berntsen, Bertrand, Chiolo, Cuckson, de Senarclens, Domiano, Doshi, Guagnano, Leblanc, Linfors, Nielsen, Pierin, Poncelet, Schell, Stergiou, Stolt, Stolt, Vinyoles).

Contributors GI and RJM conceived and designed the review. GI and JH screened the references and assessed risk of bias. GI, RS and RJM analysed the data. All authors revised and approved the final version of the review.

Funding GI is funded as an NIHR Clinical Lecturer in General Practice.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Ng M, Fleming T, Robinson M, *et al*. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384:766–81.
2. Palatini P, Parati G. Blood pressure measurement in very obese patients: a challenging problem. *J Hypertens* 2011;29:425–9.
3. Doshi H, Weder AB, Bard RL, *et al*. Does “hidden undercuffing” occur among obese patients? Effect of arm sizes and other

- predictors of the difference between wrist and upper arm blood pressures. *J Clin Hypertens (Greenwich)* 2010;12:82–8.
4. Graves JW, Bailey KR, Sheps SG. The changing distribution of arm circumferences in NHANES III and NHANES 2000 and its impact on the utility of the 'standard adult' blood pressure cuff. *Blood Press Monit* 2003;8:223–7.
 5. Graves JW. Prevalence of blood pressure cuff sizes in a referral practice of 430 consecutive adult hypertensives. *Blood Press Monit* 2001;6:17–20.
 6. National Institute of Health and Clinical Excellence. *Hypertension: clinical management of primary hypertension in adults*. London: NICE, 2011.
 7. O'Brien E. What to do when faced with an unmeasurable ambulatory blood pressure? *J Hypertens* 2011;29:451–3.
 8. Pickering TG, Hall JE, Appel LJ, et al. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension* 2005;45:142–61.
 9. Mancia G, Fagard R, Narkiewicz K, et al. 2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Blood Pressure* 2013;34:2159–219.
 10. Mendis S, Lindholm LH, Mancia G, et al. World Health Organization (WHO) and International Society of Hypertension (ISH) risk prediction charts: assessment of cardiovascular risk for prevention and control of cardiovascular disease in low and middle-income countries. *J Hypertens* 2007;25:1578–82.
 11. Lewington S, Clarke R, Qizilbash N, et al., Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002;360:1903–13.
 12. Leeflang MM, Deeks JJ, Gatsonis C, et al. Systematic reviews of diagnostic test accuracy. *Ann Intern Med* 2008;149:889–97.
 13. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011;155:529–36.
 14. Rutter CM, Gatsonis CA. A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Stat Med* 2001;20:2865–84.
 15. Altman DG, Bland JM. Commentary on quantifying agreement between two methods of measurement. *Clin Chem* 2002;48:801–2.
 16. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* 2005;58:882–93.
 17. Stergiou GS, Christodoulakis GR, Nasothimiou EG, et al. Can validated wrist devices with position sensors replace arm devices for self-home blood pressure monitoring? A randomized crossover trial using ambulatory monitoring as reference. *Am J Hypertens* 2008;21:753–8.
 18. Bertrand CA, Trout KW, Williams MH. Measurement of blood pressure in obese persons. *J Am Med Assoc* 1956;162:970–1.
 19. Bennett JM, van den Bogaerde E, van Rensburg JT. [Evaluation of a new blood pressure cuff for patients with large and normal arm circumferences]. *S Afr Med J* 1987;72:610–1.
 20. Berntsen R, Romo O, Nielsen EW. [Blood pressure measurements using three different cuff sizes]. *Tidsskr Nor Laegeforen* 1998;118:3112–15.
 21. Chiolerio A, Gervasoni JP, Rwebogora A, et al. Difference in blood pressure readings with mercury and automated devices: impact on hypertension prevalence estimates in Dar es Salaam, Tanzania. *Eur J Epidemiol* 2006;21:427–33.
 22. Cuckson AC, Moran P, Seed P, et al. Clinical evaluation of an automated oscillometric blood pressure wrist device. *Blood Press Monit* 2004;9:31–7.
 23. de Senarclens O, Feihl F, Giusti V, et al. Brachial or wrist blood pressure in obese patients: which is the best? *Blood Press Monit* 2008;13:149–51.
 24. Domiano KL, Hinck SM, Savinske DL, et al. Comparison of upper arm and forearm blood pressure. *Clin Nurs Res* 2008;17:241–50.
 25. Guagnano MT, Pace-Palitti V, Murri R, et al. The prevalence of hypertension in gynaecoid and android obese women. *J Hum Hypertens* 1996;10:619–24.
 26. Guagnano MT, Palitti VP, Murri R, et al. Many factors can affect the prevalence of hypertension in obese patients: role of cuff size and type of obesity. *Panminerva Med* 1998;40:22–7.
 27. Leblanc MÈ, Croteau S, Ferland A, et al. Blood pressure assessment in severe obesity: validation of a forearm approach. *Obesity (Silver Spring)* 2013;21:E533–41.
 28. Linfors EW, Feussner JR, Blessing CL, et al. Spurious hypertension in the obese patient. Effect of sphygmomanometer cuff size on prevalence of hypertension. *Arch Intern Med* 1984;144:1482–5.
 29. Nielsen PE, Larsen B, Holstein P, et al. Accuracy of auscultatory blood pressure measurements in hypertensive and obese subjects. *Hypertension* 1983;5:122–7.
 30. Pierin AM, Alavarce DC, Gusmão JL, et al. Blood pressure measurement in obese patients: comparison between upper arm and forearm measurements. *Blood Press Monit* 2004;9:101–5.
 31. Poncelet P, Durand P, Lechantre R, et al. [Measurement of blood pressure in obese patients: reliability and value of finger measurement (Finapres)]. *Arch Mal Coeur Vaiss* 1992;85:1165–8.
 32. Simpson JA, Jamieson G, Dickhaus DW, et al. Effect of size of cuff bladder on accuracy of measurement of indirect blood pressure. *Am Heart J* 1965;70:208–15.
 33. Stolt M, Sjonell G, Astrom H, et al. The reliability of auscultatory measurement of arterial blood pressure. A comparison of the standard and a new methodology. *Am J Hypertens* 1990;3:697–703.
 34. Stolt M, Sjonell G, Aström H, et al. Improved accuracy of indirect blood pressure measurement in patients with obese arms. *Am J Hypertens* 1993;6:66–71.
 35. Vinyoles E, Pujol E, de la Figuera M, et al. Measuring blood pressure in the forearm of obese patients: concordance with arm measurement. *Med Clin (Barc)* 2005;124:213–14.
 36. Deeks J, Bossuyt P, Gatsonis C. *Cochrane handbook for systematic reviews of diagnostic test accuracy version 1.0.0*. The Cochrane Collaboration, 2009.
 37. Blackburn H, Kihlberg J, Brozek J. Arm versus forearm blood pressure in obesity. 1965;69:423–4.
 38. Bovet P, Hungerbühler P, Quilindo J, Grettve M L, Waerber B, Burnand B. Systematic difference between blood pressure readings caused by cuff type. *Hypertension* 1994;24:786–92.
 39. Genc Y, Altunkan S, Kilinc O, Altunkan E. Comparative study on auscultatory and oscillometric methods of ambulatory blood pressure measurements in adult patients. *Blood Press Monit* 2008;13:29–35.
 40. Guagnano MT, Merlitti D, Murri R, et al. Ambulatory blood pressure monitoring in evaluating the relationship between obesity and blood pressure. *J Hum Hypertens* 1994;8:245–50.
 41. Julien J, Pagny J Y, Jeunemaitre X, Fouqueray B, Plouin P F, Corvol P. Comparison of 3 methods of blood pressure measurement in obesity. *Arch Mal Coeur Vaiss* 1988;81 Spec No:241–5.
 42. King G E. Errors in clinical measurement of blood pressure in obesity. 1967;32:223–37.
 43. Kotsis V, Stabouli S, Bouldin M, Low A, Toumanidis S, Zakopoulos N. Impact of obesity on 24-hour ambulatory blood pressure and hypertension. *Hypertension* 2005;45:602–7.
 44. Kvols L K, Rohlfing B M, Alexander J K. A comparison of intra-arterial and cuff blood pressure measurements in very obese subjects. *Cardiovasc Res Cent Bull* 1969;7:118–23.
 45. Maxwell Morton H, Schroth Phillip C, Waks Abraham U, Karam Maroun, Dornfeld Leslie P. Error in blood-pressure measurement due to incorrect cuff size in obese patients. 1982;320:33–6.
 46. Maxwell G F, Puijitt J F, Arntzenius A C. Comparison of the conical cuff and the standard rectangular cuffs. 1985;14:468–72.
 47. Pang T C, Brown M A. Accuracy of ambulatory blood pressure monitors in routine clinical practice. 2006;19:801–9.
 48. Schell K, Bradley E, Bucher L, Seckel M, Lyons D, Wakai S, et al. Clinical comparison of automatic, noninvasive measurements of blood pressure in the forearm and upper arm. *Am J Crit Care* 2005;14:232–41.
 49. Schell K, Lyons D, Bradley E, Bucher L, Seckel M, Wakai S, et al. Clinical comparison of automatic, noninvasive measurements of blood pressure in the forearm and upper arm with the patient supine or with the head of the bed raised 45 degrees: a follow-up study. *Am J Crit Care* 2006;15:196–205.
 50. Umana E, Ahmed W, Fraley M A, Alpert M A. Comparison of oscillometric and intraarterial systolic and diastolic blood pressures in lean, overweight, and obese patients. *Angiology* 2006;57:41–5.
 51. van Montfrans G, van der Hoeven G, Karemaker J, Wieling W, Dunning A. Accuracy of auscultatory blood pressure measurement with a long cuff. *Br Med J (Clin Res Ed)* 1987;295:354–5.
 52. Warembourg A, Poncelet P, Carre A. Reliability of the measurement of arterial pressure in the forearm in the obese subject. 1987;80:1015–9.
 53. Altman D, Bland M. Commentary on quantifying agreement between two methods of measurement. *Cochrane Statistical Methods Group* 2002;48:801–2.